

PARTIAL FRACTION DECOMPOSITION PROOFS OF SOME q -SERIES IDENTITIES

NANCY S. S. GU and HELMUT PRODINGER

Center for Combinatorics
LPMC-TJKLC
Nankai University
Tianjin 300071
P. R. China
e-mail: gu@nankai.edu.cn

Department of Mathematics
University of Stellenbosch
7602 Stellenbosch
South Africa

Abstract

Using the partial fraction decomposition method, we give new proofs of some q -series identities related to divisor functions in [10] and two finite q -series transformations in [8]. Likewise, some new q -series identities are obtained, including generalizations of some main results in [10] and generalizations of special cases of the q -Pfaff-Saalschütz summation theorem and the q -Chu-Vandermonde identity.

2010 Mathematics Subject Classification: 11B65, 30E05.

Keywords and phrases: q -series identity, partial fraction decomposition, divisor function.

The first author was supported by the National Natural Science Foundation of China.

The second author was supported by International Science and Technology Agreement (Grant 67215).

Received December 29, 2013

© 2013 Scientific Advances Publishers

1. Introduction

Wenchang Chu [2] showed that some seemingly difficult identities can be proved in a simple way by performing partial fraction decomposition to a suitable rational function, and then taking a certain limit. For further applications of the partial fraction decomposition method, see [1, 3-5].

Although for the cases that we study here, the method called q -Rice method [11] is computationally equivalent, we prefer to express everything in terms of partial fraction decomposition, which is conceptually simpler than contour integrals and residues.

We reprove in this way some identities from [10] and from [8] in this very simple fashion and obtain also some additional formulae. Especially, we generalize some main results in [10] and special cases of the q -Pfaff-Saalschütz summation theorem and the q -Chu-Vandermonde identity.

As usual, we follow the notation and terminology in [9]. For $|q| < 1$, the q -shifted factorial is defined by

$$(a; q)_\infty = \prod_{k=0}^{\infty} (1 - aq^k) \quad \text{and} \quad (a; q)_n = \frac{(a; q)_\infty}{(aq^n; q)_\infty}, \text{ for } n \in \mathbb{Z}.$$

For convenience, we shall adopt the following notation for multiple q -shifted factorials:

$$(a_1, a_2, \dots, a_m; q)_n = (a_1; q)_n (a_2; q)_n \cdots (a_m; q)_n,$$

where n is an integer or infinity.

The q -binomial coefficients, or the Gauss coefficients, are given by

$$\begin{bmatrix} n \\ k \end{bmatrix} = \begin{cases} \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}}, & \text{if } 0 \leq k \leq n, \\ 0, & \text{otherwise.} \end{cases}$$

The (unilateral) basic hypergeometric series ${}_r\phi_s$ is defined by

$${}_r\phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r \\ b_1, b_2, \dots, b_s \end{matrix} ; q, z \right] = \sum_{k=0}^{\infty} \frac{(a_1, a_2, \dots, a_r; q)_k}{(q, b_1, b_2, \dots, b_s; q)_k} \left[(-1)^k q^{\binom{k}{2}} \right]^{1+s-r} z^k.$$

2. Some q -Series Identities of Guo-Zhang

In [10], the authors proved new generalizations of some q -series identities of Dilcher [7] and Prodinger [11] related to divisor functions. They also obtained some special cases including an identity related to overpartitions given by Corteel and Lovejoy [6, Theorem 4.4].

In this section, using the partial fraction decomposition method, we give new proofs of some theorems in [10].

Theorem 2.1 [10, Theorem 1.1]. *For $n \geq 0$ and $0 \leq l, m \leq n$, we have*

$$\sum_{\substack{k=0 \\ k \neq m}}^n \begin{bmatrix} n \\ k \end{bmatrix} \frac{(q/t; q)_k (tq^{-l}; q)_{n-k}}{1 - q^{k-m}} t^k = (-1)^m q^{\binom{m+1}{2}} \begin{bmatrix} n \\ m \end{bmatrix} (tq^{-l}; q)_l (tq^{-m}; q)_{n-l} \\ \times \left(\sum_{k=0}^{n-l-1} \frac{tq^{k-m}}{1 - tq^{k-m}} - \sum_{\substack{k=0 \\ k \neq m}}^n \frac{q^{k-m}}{1 - q^{k-m}} \right).$$

Proof. Set

$$F(z) = \frac{(q; q)_n}{(z; q)_{n+1}} \frac{(tz; q)_{n-l} (tq^{-l}; q)_l}{(z - q^{-m})}.$$

Performing partial fraction decomposition on $F(z)$, we have

$$F(z) = \sum_{k=0}^n \frac{b_k}{1 - zq^k} + \frac{b_{n+1}}{(1 - zq^m)^2}. \quad (2.1)$$

Multiplying both sides of (2.1) by z , and then letting $z \rightarrow \infty$, we get

$$\sum_{k=0}^n \frac{b_k}{q^k} = 0. \quad (2.2)$$

Now we compute b_k for $k = 0, 1, \dots, n$.

For $k \neq m$, we have

$$\begin{aligned} b_k &= \lim_{z \rightarrow q^{-k}} (1 - zq^k) F(z) \\ &= \lim_{z \rightarrow q^{-k}} (1 - zq^k) \frac{(q; q)_n}{(z; q)_{n+1}} \frac{(tz; q)_{n-l} (tq^{-l}; q)_l}{(z - q^{-m})} \\ &= \lim_{z \rightarrow q^{-k}} \frac{(q; q)_n (tz; q)_{n-l} (tq^{-l}; q)_l}{(z; q)_k (zq^{k+1}; q)_{n-k} (z - q^{-m})} \\ &= \frac{(q; q)_n (tq^{-k}; q)_{n-l} (tq^{-l}; q)_l}{(q^{-k}; q)_k (q; q)_{n-k} (q^{-k} - q^{-m})} \\ &= \frac{(-1)^k q^{\binom{k+1}{2} + k} (q; q)_n (tq^{-k}; q)_\infty (tq^{-l}; q)_\infty}{(tq^{-k+n-l}; q)_\infty (t; q)_\infty (q; q)_k (q; q)_{n-k} (1 - q^{k-m})} \\ &= \begin{bmatrix} n \\ k \end{bmatrix} \frac{(-1)^k q^{\binom{k+1}{2} + k} (tq^{-k}; q)_k (tq^{-l}; q)_{n-k}}{1 - q^{k-m}} \\ &= \begin{bmatrix} n \\ k \end{bmatrix} \frac{(q/t; q)_k (tq^{-l}; q)_{n-k}}{1 - q^{k-m}} (tq)^k. \end{aligned}$$

For $k = m$, we have

$$\begin{aligned} b_m &= -q^{-m} \lim_{z \rightarrow q^{-m}} D_z \{(1 - zq^m)^2 F(z)\} \\ &= -q^{-m} \lim_{z \rightarrow q^{-m}} D_z \left\{ (1 - zq^m)^2 \frac{(q; q)_n}{(z; q)_{n+1}} \frac{(tz; q)_{n-l} (tq^{-l}; q)_l}{(z - q^{-m})} \right\} \end{aligned}$$

$$\begin{aligned}
&= \lim_{z \rightarrow q^{-m}} D_z \left\{ \frac{(q; q)_n (tz; q)_{n-l} (tq^{-l}; q)_l}{(z; q)_m (zq^{m+1}; q)_{n-m}} \right\} \\
&= (q; q)_n (tq^{-l}; q)_l \lim_{z \rightarrow q^{-m}} D_z \left\{ \frac{(tz; q)_{n-l}}{(z; q)_m (zq^{m+1}; q)_{n-m}} \right\} \\
&= \frac{(q; q)_n (tq^{-l}; q)_l (tq^{-m}; q)_{n-l}}{(-1)^m q^{-\binom{m+1}{2}} (q; q)_m (q; q)_{n-m}} \left(- \sum_{k=0}^{n-l-1} \frac{tq^k}{1 - tq^{k-m}} + \sum_{\substack{k=0 \\ k \neq m}}^n \frac{q^k}{1 - q^{k-m}} \right) \\
&= (-1)^{m-1} q^{\binom{m+1}{2}} \begin{bmatrix} n \\ m \end{bmatrix} (tq^{-l}; q)_l (tq^{-m}; q)_{n-l} \left(\sum_{k=0}^{n-l-1} \frac{tq^k}{1 - tq^{k-m}} - \sum_{\substack{k=0 \\ k \neq m}}^n \frac{q^k}{1 - q^{k-m}} \right).
\end{aligned}$$

Therefore, according to (2.2), we get Theorem 2.1. $\square$

Theorem 2.2 [10, Theorem 1.2]. *For $m, n \geq 1$, we have*

$$\begin{aligned}
&\sum_{k=1}^n \begin{bmatrix} n \\ k \end{bmatrix} \frac{(q^m / t; q)_k (t; q)_{n-k}}{(tq^{-m}; q)_{m+n} (1 - q^k)^m} t^k \\
&= - \sum_{1 \leq k_m \leq \dots \leq k_1 \leq n} \frac{q^{k_1}}{(1 - tq^{k_1-1})(1 - q^{k_1})} \cdots \frac{q^{k_m}}{(1 - tq^{k_m-m})(1 - q^{k_m})}.
\end{aligned}$$

Proof. Set

$$F(z) = \frac{(q; q)_n}{(z; q)_{n+1}} \frac{(tzq^{-m+1}; q)_{m+n-1}}{(tq^{-m}; q)_{m+n} (tq^{-m+1}; q)_{m-1} (z-1)^m}.$$

Performing partial fraction decomposition on $F(z)$, we have

$$F(z) = \sum_{k=0}^n \frac{b_k}{1 - zq^k} + \sum_{k=2}^{m+1} \frac{c_k}{(1 - z)^k}. \quad (2.3)$$

Multiplying both sides of (2.3) by z , and then letting $z \rightarrow \infty$, we get

$$\sum_{k=0}^n \frac{b_k}{q^k} = 0. \quad (2.4)$$

Now we calculate b_k for $0 \leq k \leq n$.

For $1 \leq k \leq n$, we have

$$\begin{aligned} b_k &= \lim_{z \rightarrow q^{-k}} (1 - zq^k) F(z) \\ &= \lim_{z \rightarrow q^{-k}} \frac{(q; q)_n (tzq^{-m+1}; q)_{m+n-1}}{(z; q)_k (zq^{k+1}; q)_{n-k} (tq^{-m}; q)_{m+n} (tq^{-m+1}; q)_{m-1} (z-1)^m} \\ &= \frac{(q; q)_n (tq^{-m-k+1}; q)_{m+n-1}}{(q^{-k}; q)_k (q; q)_{n-k} (tq^{-m}; q)_{m+n} (tq^{-m+1}; q)_{m-1} (q^{-k} - 1)^m} \\ &= \begin{bmatrix} n \\ k \end{bmatrix} \frac{(-1)^k q^{\binom{k+1}{2} + km} (tq^{-m-k+1}; q)_{m+n-1}}{(tq^{-m}; q)_{m+n} (tq^{-m+1}; q)_{m-1} (1 - q^k)^m} \\ &= \begin{bmatrix} n \\ k \end{bmatrix} \frac{(-1)^k q^{\binom{k+1}{2} + km} (tq^{-m-k+1}; q)_k (t; q)_{n-k}}{(tq^{-m}; q)_{m+n} (1 - q^k)^m} \\ &= \begin{bmatrix} n \\ k \end{bmatrix} \frac{(q^m / t; q)_k (t; q)_{n-k}}{(tq^{-m}; q)_{m+n} (1 - q^k)^m} (tq)^k. \end{aligned}$$

For $k = 0$, we have

$$\begin{aligned} b_0 &= \frac{(-1)^m}{m!} \lim_{z \rightarrow 1} D_z^{(m)} \left\{ (1 - z)^{m+1} F(z) \right\} \\ &= \frac{(-1)^m}{m!} \lim_{z \rightarrow 1} D_z^{(m)} \left\{ \frac{(-1)^m (q; q)_n (tzq^{-m+1}; q)_{m+n-1}}{(zq; q)_n (tq^{-m}; q)_{m+n} (tq^{-m+1}; q)_{m-1}} \right\} \\ &= \frac{(q; q)_n}{(tq^{-m}; q)_{m+n} (tq^{-m+1}; q)_{m-1} m!} \lim_{z \rightarrow 1} D_z^{(m)} \left\{ \frac{(tzq^{-m+1}; q)_{m+n-1}}{(zq; q)_n} \right\} \end{aligned}$$

$$\begin{aligned}
&= \frac{(q; q)_n}{(tq^{-m}; q)_m (tq^{-m+1}; q)_{m+n-1}} [(z-1)^m] \frac{(tzq^{-m+1}; q)_{m+n-1}}{(zq; q)_n} \\
&= \frac{(q; q)_n}{(tq^{-m}; q)_m (tq^{-m+1}; q)_{m+n-1}} [w^m] \frac{(t(w+1)q^{-m+1}; q)_{m+n-1}}{((w+1)q; q)_n} \\
&= \frac{1}{(tq^{-m}; q)_m} [w^m] \frac{\prod_{k=1}^{m+n-1} \left(1 - \frac{twq^{-m+k}}{1 - tq^{-m+k}}\right)}{\prod_{k=1}^n \left(1 - \frac{wq^k}{1 - q^k}\right)}.
\end{aligned}$$

In order to read off this coefficient, we have

$$\begin{aligned}
&[w^m] \frac{\prod_{k=1-m}^{n-1} \left(1 - \frac{twq^k}{1 - tq^k}\right)}{\prod_{k=1}^n \left(1 - \frac{wq^k}{1 - q^k}\right)} \\
&= \sum_{1 \leq k_m \leq \dots \leq k_1 \leq n} \left(\frac{q^{k_1}}{1 - q^{k_1}} - \frac{tq^{k_1-1}}{1 - tq^{k_1-1}} \right) \dots \left(\frac{q^{k_m}}{1 - q^{k_m}} - \frac{tq^{k_m-m}}{1 - tq^{k_m-m}} \right) \\
&= \sum_{1 \leq k_m \leq \dots \leq k_1 \leq n} \frac{q^{k_1}(1 - tq^{-1})}{(1 - tq^{k_1-1})(1 - q^{k_1})} \dots \frac{q^{k_m}(1 - tq^{-m})}{(1 - tq^{k_m-m})(1 - q^{k_m})} \\
&= (tq^{-m}; q)_m \sum_{1 \leq k_m \leq \dots \leq k_1 \leq n} \frac{q^{k_1}}{(1 - tq^{k_1-1})(1 - q^{k_1})} \dots \frac{q^{k_m}}{(1 - tq^{k_m-m})(1 - q^{k_m})}.
\end{aligned}$$

For the first equation of this chain of equalities, we argue as follows: We have to select altogether m w 's from the factors. Factors in the denominator can be chosen arbitrary often, but factors from the numerator only once. Let k_1 be the largest number such that $k_1 - 1$ from the numerator is chosen or, if not, such that k_1 is chosen from the denominator. This gives a contribution

$$\left(\frac{q^{k_1}}{1 - q^{k_1}} - \frac{tq^{k_1-1}}{1 - tq^{k_1-1}} \right).$$

Now, let $k_2 \leq k_1$ be the largest number such that $k_2 - 2$ from the numerator is chosen or, if not, such that k_2 is chosen from the denominator. This gives a contribution

$$\left(\frac{q^{k_2}}{1 - q^{k_2}} - \frac{tq^{k_2-2}}{1 - tq^{k_2-2}} \right).$$

Then, let $k_3 \leq k_2$ be the largest number such that $k_3 - 3$ from the numerator is chosen or, if not, such that k_3 is chosen from the denominator, and so on.

Therefore, we have

$$b_0 = \sum_{1 \leq k_m \leq \dots \leq k_1 \leq n} \frac{q^{k_1}}{(1 - tq^{k_1-1})(1 - q^{k_1})} \dots \frac{q^{k_m}}{(1 - tq^{k_m-m})(1 - q^{k_m})}.$$

According to (2.4), we get Theorem 2.2. □

Theorem 2.3 [10, Theorem 4.1]. *For $m, n \geq 1$, we have*

$$\begin{aligned} & \sum_{k=1}^n \left[\begin{matrix} n \\ k \end{matrix} \right] \frac{(q^m / t; q)_k (t / q; q)_{n-k}}{(tq^{-m}; q)_{m+n-1} (1 - q^k)^m} t^k \\ &= \sum_{k_1=1}^n \frac{q^{k_1}}{(1 - tq^{k_1-2})(1 - q^{k_1})} \sum_{k_2=1}^{k_1} \frac{q^{k_2}}{(1 - tq^{k_2-3})(1 - q^{k_2})} \dots \\ & \quad \times \left(\sum_{k_m=1}^{k_{m-1}-1} \frac{tq^{k_m-m}}{1 - tq^{k_m-m}} - \sum_{k_m=1}^{k_{m-1}} \frac{q^{k_m}}{1 - q^{k_m}} \right). \end{aligned}$$

Proof. Set

$$F(z) = \frac{(q; q)_n}{(z; q)_{n+1}} \frac{(tzq^{-m+1}; q)_{m+n-2}}{(tq^{-m}; q)_{m+n-1} (tq^{-m+1}; q)_{m-2} (z-1)^m}.$$

Performing partial fraction decomposition on $F(z)$, we have

$$F(z) = \sum_{k=0}^n \frac{b_k}{1 - zq^k} + \sum_{k=2}^{m+1} \frac{c_k}{(1 - z)^k}. \quad (2.5)$$

Multiplying by z on both sides of (2.5), and letting $z \rightarrow \infty$, we get

$$\sum_{k=0}^n \frac{b_k}{q^k} = 0. \quad (2.6)$$

Now we calculate b_k for $0 \leq k \leq n$.

For $1 \leq k \leq n$, we have

$$\begin{aligned} b_k &= \lim_{z \rightarrow q^{-k}} (1 - zq^k) F(z) \\ &= \lim_{z \rightarrow q^{-k}} \frac{(q; q)_n (tzq^{-m+1}; q)_{m+n-2}}{(z; q)_k (zq^{k+1}; q)_{n-k} (tq^{-m}; q)_{m+n-1} (tq^{-m+1}; q)_{m-2} (z-1)^m} \\ &= \frac{(q; q)_n (tq^{-m-k+1}; q)_{m+n-2}}{(q^{-k}; q)_k (q; q)_{n-k} (tq^{-m}; q)_{m+n-1} (tq^{-m+1}; q)_{m-2} (q^{-k} - 1)^m} \\ &= \left[\begin{matrix} n \\ k \end{matrix} \right] \frac{(-1)^k q^{\binom{k+1}{2} + km} (tq^{-m-k+1}; q)_{m+n-2}}{(tq^{-m}; q)_{m+n-1} (tq^{-m+1}; q)_{m-2} (1 - q^k)^m} \\ &= \left[\begin{matrix} n \\ k \end{matrix} \right] \frac{(-1)^k q^{\binom{k+1}{2} + km} (tq^{-m-k+1}; q)_k (t/q; q)_{n-k}}{(tq^{-m}; q)_{m+n-1} (1 - q^k)^m} \\ &= \left[\begin{matrix} n \\ k \end{matrix} \right] \frac{(q^m / t; q)_k (t/q; q)_{n-k}}{(tq^{-m}; q)_{m+n-1} (1 - q^k)^m} (tq)^k. \end{aligned}$$

For $k = 0$, we have

$$b_0 = \frac{(-1)^m}{m!} \lim_{z \rightarrow 1} D_z^{(m)} \left\{ (1 - z)^{m+1} F(z) \right\}$$

$$\begin{aligned}
&= \frac{(-1)^m}{m!} \lim_{z \rightarrow 1} D_z^{(m)} \left\{ \frac{(-1)^m (q; q)_n (tzq^{-m+1}; q)_{m+n-2}}{(zq; q)_n (tq^{-m}; q)_{m+n-1} (tq^{-m+1}; q)_{m-2}} \right\} \\
&= \frac{(q; q)_n}{(tq^{-m}; q)_{m+n-1} (tq^{-m+1}; q)_{m-2} m!} \lim_{z \rightarrow 1} D_z^{(m)} \left\{ \frac{(tzq^{-m+1}; q)_{m+n-2}}{(zq; q)_n} \right\} \\
&= \frac{(q; q)_n}{(tq^{-m}; q)_{m+n-1} (tq^{-m+1}; q)_{m-2}} [(z-1)^m] \frac{(tzq^{-m+1}; q)_{m+n-2}}{(zq; q)_n} \\
&= \frac{(q; q)_n}{(tq^{-m}; q)_{m+n-1} (tq^{-m+1}; q)_{m-2}} [w^m] \frac{(t(w+1)q^{-m+1}; q)_{m+n-2}}{((w+1)q; q)_n} \\
&= \frac{1}{(tq^{-m}; q)_{m-1}} [w^m] \frac{\prod_{k=1}^{m+n-2} \left(1 - \frac{twq^{-m+k}}{1 - tq^{-m+k}} \right)}{\prod_{k=1}^n \left(1 - \frac{wq^k}{1 - q^k} \right)}.
\end{aligned}$$

In order to read off this coefficient, we have

$$\begin{aligned}
&[w^m] \frac{\prod_{k=1-m}^{n-2} \left(1 - \frac{twq^k}{1 - tq^k} \right)}{\prod_{k=1}^n \left(1 - \frac{wq^k}{1 - q^k} \right)} \\
&= \sum_{k_1=1}^n \left(\frac{q^{k_1}}{1 - q^{k_1}} - \frac{tq^{k_1-2}}{1 - tq^{k_1-2}} \right) \sum_{k_2=1}^{k_1} \left(\frac{q^{k_2}}{1 - q^{k_2}} - \frac{tq^{k_2-3}}{1 - tq^{k_2-3}} \right) \dots \\
&\quad \times \sum_{k_{m-1}=1}^{k_{m-2}} \left(\frac{q^{k_{m-1}}}{1 - q^{k_{m-1}}} - \frac{tq^{k_{m-1}-m}}{1 - tq^{k_{m-1}-m}} \right) \\
&\quad \times \left[\sum_{k_m=1}^{k_{m-1}} \left(\frac{q^{k_m}}{1 - q^{k_m}} - \frac{tq^{k_m-m-1}}{1 - tq^{k_m-m-1}} \right) - \left(-\frac{tq^{-m}}{1 - tq^{-m}} \right) \right]
\end{aligned}$$

$$\begin{aligned}
&= \sum_{k_1=1}^n \frac{q^{k_1}(1-tq^{-2})}{(1-tq^{k_1-2})(1-q^{k_1})} \sum_{k_2=1}^{k_1} \frac{q^{k_2}(1-tq^{-3})}{(1-tq^{k_2-3})(1-q^{k_2})} \cdots \\
&\quad \times \sum_{k_{m-1}=1}^{k_{m-2}} \frac{q^{k_{m-1}}(1-tq^{-m})}{(1-tq^{k_{m-1}-m})(1-q^{k_{m-1}})} \\
&\quad \times \left(\sum_{k_m=1}^{k_{m-1}} \frac{q^{k_m}}{1-q^{k_m}} - \sum_{k_m=1}^{k_{m-1}} \frac{tq^{k_m-m-1}}{1-tq^{k_m-m-1}} + \frac{tq^{-m}}{1-tq^{-m}} \right) \\
&= (tq^{-m}; q)_{m-1} \sum_{k_1=1}^n \frac{q^{k_1}}{(1-tq^{k_1-2})(1-q^{k_1})} \sum_{k_2=1}^{k_1} \frac{q^{k_2}}{(1-tq^{k_2-3})(1-q^{k_2})} \cdots \\
&\quad \times \sum_{k_{m-1}=1}^{k_{m-2}} \frac{q^{k_{m-1}}}{(1-tq^{k_{m-1}-m})(1-q^{k_{m-1}})} \left(\sum_{k_m=1}^{k_{m-1}} \frac{q^{k_m}}{1-q^{k_m}} - \sum_{k_m=2}^{k_{m-1}} \frac{tq^{k_m-m-1}}{1-tq^{k_m-m-1}} \right) \\
&= (tq^{-m}; q)_{m-1} \sum_{k_1=1}^n \frac{q^{k_1}}{(1-tq^{k_1-2})(1-q^{k_1})} \sum_{k_2=1}^{k_1} \frac{q^{k_2}}{(1-tq^{k_2-3})(1-q^{k_2})} \cdots \\
&\quad \times \sum_{k_{m-1}=1}^{k_{m-2}} \frac{q^{k_{m-1}}}{(1-tq^{k_{m-1}-m})(1-q^{k_{m-1}})} \left(\sum_{k_m=1}^{k_{m-1}} \frac{q^{k_m}}{1-q^{k_m}} - \sum_{k_m=1}^{k_{m-1}-1} \frac{tq^{k_m-m}}{1-tq^{k_m-m}} \right).
\end{aligned}$$

The explanation is very similar to the earlier instance Theorem 2.2. The difference is that, if $k_m = 1$, there is an exception, since $k_m - m - 1 = -m$ is out of range for the numerator, and thus such a term cannot be taken.

Perhaps a more aesthetic way would be to write

$$\begin{aligned}
&[w^m] \frac{\prod_{k=1-m}^{n-2} \left(1 - \frac{twq^k}{1-tq^k} \right)}{\prod_{k=1}^n \left(1 - \frac{wq^k}{1-q^k} \right)} \\
&= \sum_{2 \leq k_m \leq \cdots \leq k_1 \leq n} \left(\frac{q^{k_1}}{1-q^{k_1}} - \frac{tq^{k_1-2}}{1-tq^{k_1-2}} \right) \cdots \left(\frac{q^{k_m}}{1-q^{k_m}} - \frac{tq^{k_m-m-1}}{1-tq^{k_m-m-1}} \right)
\end{aligned}$$

$$\begin{aligned}
& + \sum_{1 \leq k_{m-1} \leq \dots \leq k_1 \leq n} \left(\frac{q^{k_1}}{1 - q^{k_1}} - \frac{tq^{k_1-2}}{1 - tq^{k_1-2}} \right) \dots \left(\frac{q^{k_{m-1}}}{1 - q^{k_{m-1}}} - \frac{tq^{k_{m-1}-m}}{1 - tq^{k_{m-1}-m}} \right) \frac{q}{1 - q} \\
& = \sum_{2 \leq k_m \leq \dots \leq k_1 \leq n} \frac{q^{k_1}(1 - tq^{-2})}{(1 - q^{k_1})(1 - tq^{k_1-2})} \dots \frac{q^{k_m}(1 - tq^{-m-1})}{(1 - q^{k_m})(1 - tq^{k_m-m-1})} \\
& + \sum_{1 \leq k_{m-1} \leq \dots \leq k_1 \leq n} \frac{q^{k_1}(1 - tq^{-2})}{(1 - q^{k_1})(1 - tq^{k_1-2})} \dots \frac{q^{k_{m-1}}(1 - tq^{-m})}{(1 - q^{k_{m-1}})(1 - tq^{k_{m-1}-m})} \frac{q}{1 - q} \\
& = (tq^{-m-1}; q)_m \sum_{2 \leq k_m \leq \dots \leq k_1 \leq n} \frac{q^{k_1}}{(1 - q^{k_1})(1 - tq^{k_1-2})} \dots \frac{q^{k_m}}{(1 - q^{k_m})(1 - tq^{k_m-m-1})} \\
& + (tq^{-m}; q)_{m-1} \sum_{1 \leq k_{m-1} \leq \dots \leq k_1 \leq n} \frac{q^{k_1}}{(1 - q^{k_1})(1 - tq^{k_1-2})} \dots \frac{q^{k_{m-1}}}{(1 - q^{k_{m-1}})(1 - tq^{k_{m-1}-m})} \frac{q}{1 - q}.
\end{aligned}$$

Therefore, we have

$$\begin{aligned}
b_0 & = \sum_{k_1=1}^n \frac{q^{k_1}}{(1 - tq^{k_1-2})(1 - q^{k_1})} \sum_{k_2=1}^{k_1} \frac{q^{k_2}}{(1 - tq^{k_2-3})(1 - q^{k_2})} \dots \\
& \quad \times \left(\sum_{k_m=1}^{k_{m-1}} \frac{q^{k_m}}{1 - q^{k_m}} - \sum_{k_m=1}^{k_{m-1}-1} \frac{tq^{k_m-m}}{1 - tq^{k_m-m}} \right).
\end{aligned}$$

According to (2.6), we get Theorem 2.3. $\square$

3. New Results

In this section, using the partial fraction decomposition method, we get new generalizations of some main results in [10], and also obtain some other new q -series identities, including generalizations of special cases of the q -Pfaff-Saalschütz summation theorem and the q -Chu-Vandermonde identity.

After that, we give proofs of three theorems given by Guo and Zhang in [10]. While we do not find a proper way to prove [10, Theorem 1.3] by using the partial fraction decomposition method, we do get the following similar new result:

Theorem 3.1. *For $m, n \geq 1$, we have*

$$\begin{aligned} & \sum_{k=1}^n \frac{(q/t; q)_k (vq^m; q)_k (t; q)_{n-k} (t; q)_m}{(q; q)_k (v; q)_k (q; q)_{n-k} (q^k; q)_{m+1}} t^k \\ & \quad - \sum_{k=1}^m \frac{(tq^k; q)_{m-k} (t; q)_{n+k} (q/v; q)_k}{(q^k; q)_{n+1} (vq^{m-k}; q)_k (q; q)_{m-k} (q; q)_k} v^k \\ & = \frac{(t; q)_m (t; q)_n}{(q; q)_m (q; q)_n} \left(\sum_{k=1}^m \frac{q^k (1-v/q)}{(1-vq^{k-1})(1-q^k)} - \sum_{k=1}^n \frac{q^k (1-t/q)}{(1-tq^{k-1})(1-q^k)} \right). \end{aligned} \quad (3.1)$$

Proof. Set

$$F(z) = - \frac{(tz; q)_n (t; q)_m (zq^{1-m}/v; q)_m}{(z; q)_{n+1} (v; q)_m (zq^{-m}; q)_{m+1}} \left(\frac{v}{q} \right)^m.$$

Performing partial fraction decomposition on $F(z)$, we have

$$F(z) = \sum_{k=1}^n \frac{b_k}{1-zq^k} + \sum_{k=1}^m \frac{c_k}{1-zq^{-k}} + \frac{a_1}{1-z} + \frac{a_2}{(1-z)^2}. \quad (3.2)$$

Multiplying by z on both sides of (3.2), and letting $z \rightarrow \infty$, we get

$$\sum_{k=1}^n \frac{b_k}{q^k} + \sum_{k=1}^m \frac{c_k}{q^{-k}} = -a_1. \quad (3.3)$$

For $1 \leq k \leq n$, we have

$$\begin{aligned} b_k &= \lim_{z \rightarrow q^{-k}} (1-zq^k) F(z) \\ &= - \lim_{z \rightarrow q^{-k}} \frac{(tz; q)_n (t; q)_m (zq^{1-m}/v; q)_m}{(z; q)_k (zq^{k+1}; q)_{n-k} (v; q)_m (zq^{-m}; q)_{m+1}} \left(\frac{v}{q} \right)^m \end{aligned}$$

$$\begin{aligned}
&= -\frac{(tq^{-k}; q)_n(t; q)_m(q^{-m-k+1}/v; q)_m}{(q^{-k}; q)_k(q; q)_{n-k}(v; q)_m(q^{-m-k}; q)_{m+1}} \left(\frac{v}{q}\right)^m \\
&= \frac{(-1)^k q^{\binom{k+1}{2}+k} (tq^{-k}; q)_n(t; q)_m(vq^k; q)_m}{(q; q)_k(q; q)_{n-k}(v; q)_m(q^k; q)_{m+1}} \\
&= \frac{(-1)^k q^{\binom{k+1}{2}+k} (tq^{-k}; q)_k(t; q)_m(t; q)_{n-k}(vq^m; q)_k}{(q; q)_k(q; q)_{n-k}(v; q)_k(q^k; q)_{m+1}} \\
&= \frac{(q/t; q)_k(vq^m; q)_k(t; q)_{n-k}(t; q)_m}{(q; q)_k(v; q)_k(q; q)_{n-k}(q^k; q)_{m+1}} (tq)^k.
\end{aligned}$$

For $1 \leq k \leq m$, we have

$$\begin{aligned}
c_k &= \lim_{z \rightarrow q^k} (1 - zq^{-k})F(z) \\
&= -\lim_{z \rightarrow q^k} \frac{(tz; q)_n(t; q)_m(zq^{1-m}/v; q)_m}{(z; q)_{n+1}(v; q)_m(zq^{-m}; q)_{m-k}(zq^{-k+1}; q)_k} \left(\frac{v}{q}\right)^m \\
&= -\frac{(tq^k; q)_n(t; q)_m(q^{-m+k+1}/v; q)_m}{(q^k; q)_{n+1}(v; q)_m(q^{-m+k}; q)_{m-k}(q; q)_k} \left(\frac{v}{q}\right)^m \\
&= \frac{(-1)^{k+1} q^{\binom{k}{2}} (tq^k; q)_n(t; q)_m(vq^{-k}; q)_m}{(q^k; q)_{n+1}(v; q)_m(q; q)_{m-k}(q; q)_k} \\
&= \frac{(-1)^{k+1} q^{\binom{k}{2}} (tq^k; q)_{m-k}(t; q)_{n+k}(vq^{-k}; q)_k}{(q^k; q)_{n+1}(vq^{m-k}; q)_k(q; q)_{m-k}(q; q)_k} \\
&= -\frac{(tq^k; q)_{m-k}(t; q)_{n+k}(q/v; q)_k}{(q^k; q)_{n+1}(vq^{m-k}; q)_k(q; q)_{m-k}(q; q)_k} \left(\frac{v}{q}\right)^k.
\end{aligned}$$

Now we compute a_1

$$\begin{aligned}
a_1 &= -\lim_{z \rightarrow 1} D_z \{(1-z)^2 F(z)\} \\
&= \lim_{z \rightarrow 1} D_z \left\{ \frac{(tz; q)_n (t; q)_m (zq^{1-m} / v; q)_m \left(\frac{v}{q}\right)^m}{(zq; q)_n (v; q)_m (zq^{-m}; q)_m} \right\} \\
&= \frac{(t; q)_m \left(\frac{v}{q}\right)^m}{(v; q)_m} \lim_{z \rightarrow 1} D_z \left\{ \frac{(tz; q)_n (zq^{1-m} / v; q)_m}{(zq; q)_n (zq^{-m}; q)_m} \right\} \\
&= \frac{(t; q)_m \left(\frac{v}{q}\right)^m}{(v; q)_m} \lim_{z \rightarrow 1} \left[\frac{(tz; q)_n (zq^{1-m} / v; q)_m}{(zq; q)_n (zq^{-m}; q)_m} \right. \\
&\quad \times \left(-\sum_{k=0}^{n-1} \frac{tq^k}{1-tzq^k} + \sum_{k=0}^{m-1} \frac{1}{z-vq^k} + \sum_{k=1}^n \frac{q^k}{1-zq^k} - \sum_{k=1}^m \frac{1}{z-q^k} \right) \Big] \\
&= \frac{(t; q)_m (t; q)_n (q^{1-m} / v; q)_m \left(\frac{v}{q}\right)^m}{(v; q)_m (q; q)_n (q^{-m}; q)_m} \\
&\quad \times \left(-\sum_{k=0}^{n-1} \frac{tq^k}{1-tq^k} + \sum_{k=0}^{m-1} \frac{1}{1-vq^k} + \sum_{k=1}^n \frac{q^k}{1-q^k} - \sum_{k=1}^m \frac{1}{1-q^k} \right) \\
&= \frac{(t; q)_m (t; q)_n}{(q; q)_m (q; q)_n} \left[\sum_{k=1}^n \left(\frac{q^k}{1-q^k} - \frac{tq^{k-1}}{1-tq^{k-1}} \right) + \sum_{k=1}^m \left(\frac{1}{1-vq^{k-1}} - \frac{1}{1-q^k} \right) \right] \\
&= \frac{(t; q)_m (t; q)_n}{(q; q)_m (q; q)_n} \left(\sum_{k=1}^n \frac{q^k (1-t/q)}{(1-tq^{k-1})(1-q^k)} - \sum_{k=1}^m \frac{q^k (1-v/q)}{(1-vq^{k-1})(1-q^k)} \right).
\end{aligned}$$

According to (3.3), we get (3.1). $\square$

When we set $v = t$ in (3.1), we obtain

$$\sum_{k=1}^n \frac{(t; q)_{n-k} (t; q)_{m+k} (q/t; q)_k}{(q^k; q)_{m+1} (q; q)_{n-k} (q; q)_k (t; q)_k} t^k - \sum_{k=1}^m \frac{(t; q)_{m-k} (t; q)_{n+k} (q/t; q)_k}{(q^k; q)_{n+1} (q; q)_{m-k} (q; q)_k (t; q)_k} t^k$$

$$= \frac{(1-t/q)(t; q)_m(t; q)_n}{(q; q)_m(q; q)_n} \left(\sum_{k=1}^m \frac{q^k}{(1-tq^{k-1})(1-q^k)} - \sum_{k=1}^n \frac{q^k}{(1-tq^{k-1})(1-q^k)} \right), \quad (3.4)$$

which is $v = t$ case of the following theorem given by Guo and Zhang in [10].

Theorem 3.2 [10, Theorem 1.3]. *For $m, n \geq 0$, we have*

$$\begin{aligned} & \sum_{k=1}^n \frac{(q/t; q)_k(vq^m; q)_k(t; q)_{n-k}(t; q)_m}{(q; q)_k(v; q)_k(q; q)_{n-k}(q^k; q)_{m+1}} t^k - \sum_{k=1}^m \frac{(q/t; q)_k(vq^n; q)_k(t; q)_{m-k}(t; q)_n}{(q; q)_k(v; q)_k(q; q)_{m-k}(q^k; q)_{n+1}} t^k \\ &= \frac{(1-t/q)(t; q)_m(t; q)_n}{(q; q)_m(q; q)_n} \left(\sum_{k=1}^m \frac{q^k}{(1-tq^{k-1})(1-q^k)} - \sum_{k=1}^n \frac{q^k}{(1-tq^{k-1})(1-q^k)} \right). \end{aligned}$$

Interchanging m, n in (3.1), we have

$$\begin{aligned} & \sum_{k=1}^m \frac{(q/t; q)_k(vq^n; q)_k(t; q)_{m-k}(t; q)_n}{(q; q)_k(v; q)_k(q; q)_{m-k}(q^k; q)_{n+1}} t^k \\ & \quad - \sum_{k=1}^n \frac{(tq^k; q)_{n-k}(t; q)_{m+k}(q/v; q)_k}{(q^k; q)_{m+1}(vq^{n-k}; q)_k(q; q)_{n-k}(q; q)_k} v^k \\ &= \frac{(t; q)_m(t; q)_n}{(q; q)_m(q; q)_n} \left(\sum_{k=1}^n \frac{q^k(1-v/q)}{(1-vq^{k-1})(1-q^k)} - \sum_{k=1}^m \frac{q^k(1-t/q)}{(1-tq^{k-1})(1-q^k)} \right). \end{aligned} \quad (3.5)$$

Combining (3.1) and (3.5), we get

$$\begin{aligned} & \sum_{k=1}^n \frac{(q/t; q)_k(vq^m; q)_k(t; q)_{n-k}(t; q)_m}{(q; q)_k(v; q)_k(q; q)_{n-k}(q^k; q)_{m+1}} t^k \\ & \quad - \sum_{k=1}^m \frac{(q/t; q)_k(vq^n; q)_k(t; q)_{m-k}(t; q)_n}{(q; q)_k(v; q)_k(q; q)_{m-k}(q^k; q)_{n+1}} t^k \end{aligned}$$

$$\begin{aligned}
&= \sum_{k=1}^m \frac{(tq^k; q)_{m-k}(t; q)_{n+k}(q/v; q)_k}{(q^k; q)_{n+1}(vq^{m-k}; q)_k(q; q)_{m-k}(q; q)_k} v^k \\
&\quad - \sum_{k=1}^n \frac{(tq^k; q)_{n-k}(t; q)_{m+k}(q/v; q)_k}{(q^k; q)_{m+1}(vq^{n-k}; q)_k(q; q)_{n-k}(q; q)_k} v^k \\
&\quad + \frac{(1-v/q)(t; q)_m(t; q)_n}{(q; q)_m(q; q)_n} \left(\sum_{k=1}^m \frac{q^k}{(1-vq^{k-1})(1-q^k)} - \sum_{k=1}^n \frac{q^k}{(1-vq^{k-1})(1-q^k)} \right) \\
&\quad + \frac{(1-t/q)(t; q)_m(t; q)_n}{(q; q)_m(q; q)_n} \left(\sum_{k=1}^m \frac{q^k}{(1-tq^{k-1})(1-q^k)} - \sum_{k=1}^n \frac{q^k}{(1-tq^{k-1})(1-q^k)} \right).
\end{aligned}$$

According to the above identity and Theorem 3.2, we have the following identity:

$$\begin{aligned}
&\sum_{k=1}^n \frac{(tq^k; q)_{n-k}(t; q)_{m+k}(q/v; q)_k}{(q^k; q)_{m+1}(vq^{n-k}; q)_k(q; q)_{n-k}(q; q)_k} v^k \\
&\quad - \sum_{k=1}^m \frac{(tq^k; q)_{m-k}(t; q)_{n+k}(q/v; q)_k}{(q^k; q)_{n+1}(vq^{m-k}; q)_k(q; q)_{m-k}(q; q)_k} v^k \\
&= \frac{(1-v/q)(t; q)_m(t; q)_n}{(q; q)_m(q; q)_n} \left(\sum_{k=1}^m \frac{q^k}{(1-vq^{k-1})(1-q^k)} - \sum_{k=1}^n \frac{q^k}{(1-vq^{k-1})(1-q^k)} \right),
\end{aligned}$$

which can be obtained by interchanging v and t in Theorem 3.2.

Remark 3.3. We were not successful to prove [10, Theorem 1.3] with the partial fraction decomposition method. However, we offer the following observation. The right hand-side of it does not depend on the parameter v . If this fact could be shown by simple means, then we could argue that the left hand-side also does not depend on v . Henceforth, we could set $v = t$, and would achieve a proof.

Next, making some changes on the rational function $F(z)$ in the proof of Theorem 2.1, we can get some new results.

Set

$$F(z) = \frac{(vq; q)_n(tz; q)_{n-l}(tq^{-l}/v; q)_l}{(zv; q)_{n+1}(z - q^{-m})}.$$

Then using the partial fraction decomposition method, we get

$$\sum_{k=0}^n \frac{(vq/t; q)_k(tq^{-l}/v; q)_{n-k} \left(\frac{t}{v}\right)^k}{(q; q)_k(q; q)_{n-k}(1 - vq^{k-m})} = \frac{(tq^{-m}; q)_{n-l}(tq^{-l}/v; q)_l}{(vq^{-m}; q)_{n+1}}. \quad (3.6)$$

Setting v and t to be aq^m and aq^{m+1}/b in (3.6), respectively, we get the following result:

Theorem 3.4. *For $n \geq 0$ and $0 \leq l \leq n$, we have*

$$\sum_{k=0}^n \frac{(a, b, q^{-n}; q)_k}{(q, aq, bq^{-n+l}; q)_k} (q^{l+1})^k = \frac{(q; q)_n(aq/b; q)_{n-l}}{(aq; q)_n(q/b; q)_{n-l}}. \quad (3.7)$$

Setting $l = 0$ in (3.7), we obtain

$$\sum_{k=0}^n \frac{(a, b, q^{-n}; q)_k}{(q, aq, bq^{-n}; q)_k} q^k = \frac{(q, aq/b; q)_n}{(aq, q/b; q)_n},$$

which is $c = aq$ case of the q -Pfaff-Saalschütz summation theorem [9, Appendix II.12]

$${}_3\phi_2 \left[\begin{matrix} q^{-n}, & a, & b \\ & c, & abq^{1-n}/c \end{matrix} ; q, q \right] = \frac{(c/a, c/b; q)_n}{(c, c/ab; q)_n}.$$

If we set $a = q$ in (3.7), we get

Corollary 3.5. *For $n \geq 0$ and $0 \leq l \leq n$, we have*

$$\sum_{k=0}^n \frac{(b, q^{-n}; q)_k}{(q^2, bq^{-n+l}; q)_k} (q^{l+1})^k = \frac{(q; q)_n(q^2/b; q)_{n-l}}{(q^2; q)_n(q/b; q)_{n-l}}. \quad (3.8)$$

When we set $b = q^{n-l+1}$ in (3.8), we get

$$\sum_{k=0}^n \frac{(q^{n-l+1}, q^{-n}; q)_k}{(q, q^2; q)_k} (q^{l+1})^k = 0,$$

which is $a = q^{n-l+1}$ and $c = q^2$ case of the q -Chu-Vandermonde identity [9, Appendix II.7]

$${}_2\phi_1 \left[\begin{matrix} q^{-n}, & a \\ & c \end{matrix} ; q, \frac{cq^n}{a} \right] = \frac{(c/a; q)_n}{(c; q)_n}.$$

Moreover, we find that if we put a new parameter v in the proofs of Theorems 2.2 and 2.3, we get generalizations of these two theorems.

First, we give a generalization of Theorem 2.2.

Theorem 3.6. *For $m, n \geq 1$, we have*

$$\begin{aligned} & \sum_{k=1}^n \frac{(vq; q)_n (vq^{m+1}/t; q)_{k-1} (t/v; q)_{n-k}}{(tq^{-m+1}; q)_{m+n-1} (q; q)_{k-1} (q; q)_{n-k} (1-vq^k)^{m+1}} \left(\frac{t}{v}\right)^{k-1} \\ &= \sum_{1 \leq k_m \leq \dots \leq k_1 \leq n} \frac{q^{k_1-1}}{(1-tq^{k_1-1})(1-vq^{k_1})} \cdots \frac{q^{k_m-1}}{(1-tq^{k_m-m})(1-vq^{k_m})}. \end{aligned}$$

Proof. Set

$$F(z) = \frac{(vq; q)_n (tzq^{-m+1}; q)_{m+n-1}}{(vzq; q)_n (tq^{-m}/v; q)_m (tq^{-m+1}; q)_{m+n-1} (z-1)^{m+1}}.$$

Performing partial fraction decomposition on $F(z)$, we have

$$F(z) = \sum_{k=1}^n \frac{b_k}{1-vzq^k} + \sum_{k=1}^{m+1} \frac{c_k}{(1-z)^k}. \quad (3.9)$$

Multiplying both sides of (3.9) by z , and then letting $z \rightarrow \infty$, we get

$$\sum_{k=1}^n \frac{b_k}{vq^k} = -c_1. \quad (3.10)$$

Now we calculate b_k for $1 \leq k \leq n$.

For $1 \leq k \leq n$, we have

$$\begin{aligned}
b_k &= \lim_{z \rightarrow q^{-k}/v} (1 - v z q^k) F(z) \\
&= \lim_{z \rightarrow q^{-k}/v} \frac{(vq; q)_n (tzq^{-m+1}; q)_{m+n-1}}{(vzq; q)_{k-1} (vzq^{k+1}; q)_{n-k} (tq^{-m}/v; q)_m (tq^{-m+1}; q)_{m+n-1} (z-1)^{m+1}} \\
&= \frac{(vq; q)_n (tq^{-m-k+1}/v; q)_{m+n-1}}{(q^{1-k}; q)_{k-1} (q; q)_{n-k} (tq^{-m}/v; q)_m (tq^{-m+1}; q)_{m+n-1} (q^{-k}/v - 1)^{m+1}} \\
&= \frac{(-1)^{k-1} q^{\binom{k+1}{2} + km} v^{m+1} (vq; q)_n (tq^{-m-k+1}/v; q)_{m+n-1}}{(q; q)_{k-1} (q; q)_{n-k} (tq^{-m}/v; q)_m (tq^{-m+1}; q)_{m+n-1} (1 - vq^k)^{m+1}} \\
&= \frac{(-1)^{k-1} q^{\binom{k+1}{2} + km} v^{m+1} (vq; q)_n (tq^{-m-k+1}/v; q)_{k-1} (t/v; q)_{n-k}}{(q; q)_{k-1} (q; q)_{n-k} (tq^{-m+1}; q)_{m+n-1} (1 - vq^k)^{m+1}} \\
&= \frac{(vq; q)_n (vq^{m+1}/t; q)_{k-1} (t/v; q)_{n-k}}{(q; q)_{k-1} (q; q)_{n-k} (tq^{-m+1}; q)_{m+n-1} (1 - vq^k)^{m+1}} \left(\frac{t}{v}\right)^{k-1} v^{m+1} q^{m+k}.
\end{aligned}$$

For c_1 , we have

$$\begin{aligned}
c_1 &= \frac{(-1)^m}{m!} \lim_{z \rightarrow 1} D_z^{(m)} \{(1-z)^{m+1} F(z)\} \\
&= \frac{(-1)^m}{m!} \lim_{z \rightarrow 1} D_z^{(m)} \left\{ \frac{(-1)^{m+1} (vq; q)_n (tzq^{-m+1}; q)_{m+n-1}}{(vzq; q)_n (tq^{-m}/v; q)_m (tq^{-m+1}; q)_{m+n-1}} \right\} \\
&= - \frac{(vq; q)_n}{(tq^{-m}/v; q)_m (tq^{-m+1}; q)_{m+n-1}} \lim_{z \rightarrow 1} D_z^{(m)} \left\{ \frac{(tzq^{-m+1}; q)_{m+n-1}}{(vzq; q)_n} \right\} \\
&= - \frac{(vq; q)_n}{(tq^{-m}/v; q)_m (tq^{-m+1}; q)_{m+n-1}} [(z-1)^m] \frac{(tzq^{-m+1}; q)_{m+n-1}}{(vzq; q)_n}
\end{aligned}$$

$$\begin{aligned}
&= - \frac{(vq; q)_n}{(tq^{-m}/v; q)_m (tq^{-m+1}; q)_{m+n-1}} [w^m] \frac{(t(w+1)q^{-m+1}; q)_{m+n-1}}{(v(w+1)q; q)_n} \\
&= - \frac{1}{(tq^{-m}/v; q)_m} [w^m] \frac{\prod_{k=1}^{m+n-1} \left(1 - \frac{twq^{-m+k}}{1 - tq^{-m+k}}\right)}{\prod_{k=1}^n \left(1 - \frac{vwq^k}{1 - vq^k}\right)} \\
&= - \frac{1}{(tq^{-m}/v; q)_m} \sum_{1 \leq k_m \leq \dots \leq k_1 \leq n} \left(\frac{vq^{k_1}}{1 - vq^{k_1}} - \frac{tq^{k_1-1}}{1 - tq^{k_1-1}} \right) \\
&\quad \dots \left(\frac{vq^{k_m}}{1 - vq^{k_m}} - \frac{tq^{k_m-m}}{1 - tq^{k_m-m}} \right) \\
&= - \frac{1}{(tq^{-m}/v; q)_m} \sum_{1 \leq k_m \leq \dots \leq k_1 \leq n} \frac{vq^{k_1}(1 - tq^{-1}/v)}{(1 - tq^{k_1-1})(1 - vq^{k_1})} \\
&\quad \dots \frac{vq^{k_m}(1 - tq^{-m}/v)}{(1 - tq^{k_m-m})(1 - vq^{k_m})} \\
&= - v^m \sum_{1 \leq k_m \leq \dots \leq k_1 \leq n} \frac{q^{k_1}}{(1 - tq^{k_1-1})(1 - vq^{k_1})} \dots \frac{q^{k_m}}{(1 - tq^{k_m-m})(1 - vq^{k_m})}.
\end{aligned}$$

We read off the coefficient of w^m as we did in the proof of Theorem 2.2.

According to (3.10), we get Theorem 3.6. $\square$

Setting $v = 1$ in Theorem 3.6, we get Theorem 2.2.

Setting $v = 0$ in Theorem 3.6, we have

$$\sum_{k=1}^n \begin{bmatrix} n-1 \\ k-1 \end{bmatrix} (-1)^k q^{\binom{k+1}{2} - nk} = 0,$$

which is a special case of the q -binomial theorem [9, Appendix II.4]

$${}_1\phi_0 \left[\begin{matrix} q^{-n} \\ - \end{matrix} ; q, z \right] = (zq^{-n}; q)_n.$$

Setting $v = q$ in Theorem 3.6, we have the following result:

Corollary 3.7. *For $m, n \geq 1$, we have*

$$\begin{aligned} & \sum_{k=1}^n \frac{(q^2; q)_n (q^{m+2} / t; q)_{k-1} (t / q; q)_{n-k}}{(tq^{-m+1}; q)_{m+n-1} (q; q)_{k-1} (q; q)_{n-k} (1 - q^{k+1})^{m+1}} \left(\frac{t}{q} \right)^{k-1} \\ &= \sum_{1 \leq k_m \leq \dots \leq k_1 \leq n} \frac{q^{k_1-1}}{(1 - tq^{k_1-1})(1 - q^{k_1+1})} \cdots \frac{q^{k_m-1}}{(1 - tq^{k_m-m})(1 - q^{k_m+1})}. \end{aligned}$$

Letting $n \rightarrow \infty$ in Theorem 3.6, we have

Corollary 3.8. *For $m \geq 1$, we have*

$$\begin{aligned} & \sum_{k=1}^{\infty} \frac{(vq^{m+1} / t; q)_{k-1}}{(tq^{-m+1}; q)_{m-1} (q; q)_{k-1} (1 - vq^k)^{m+1}} \left(\frac{t}{v} \right)^{k-1} \\ &= \frac{(q; q)_{\infty} (t; q)_{\infty}}{(vq; q)_{\infty} (t / v; q)_{\infty}} \sum_{1 \leq k_m \leq \dots \leq k_1 < \infty} \frac{q^{k_1-1}}{(1 - tq^{k_1-1})(1 - vq^{k_1})} \\ & \quad \cdots \frac{q^{k_m-1}}{(1 - tq^{k_m-m})(1 - vq^{k_m})}. \end{aligned}$$

Setting $m = 1$ in Theorem 3.6, we get the following generalization of [10, Corollary 3.3].

Corollary 3.9. *For $n \geq 1$, we have*

$$\sum_{k=1}^n \frac{(vq; q)_n (vq^2 / t; q)_{k-1} (t / v; q)_{n-k}}{(t, q)_n (q; q)_{k-1} (q; q)_{n-k} (1 - vq^{k_m})} \left(\frac{t}{v} \right)^{k-1} = \sum_{k=1}^n \frac{q^{k-1}}{(1 - tq^{k-1})(1 - vq^k)}.$$

Now we give the following theorem, which is a generalization of Theorem 2.3.

Theorem 3.10. *For $m, n \geq 1$, we have*

$$\begin{aligned} & \sum_{k=1}^n \frac{(vq; q)_n (vq^{m+1} / t; q)_{k-1} (tq^{-1} / v; q)_{n-k}}{(tq^{-m+1}; q)_{m+n-2} (q; q)_{k-1} (q; q)_{n-k} (1 - vq^k)^{m+1}} \left(\frac{t}{v} \right)^{k-1} \\ &= \sum_{k_1=1}^n \frac{q^{k_1-1}}{(1 - tq^{k_1-2}) (1 - vq^{k_1})} \cdots \sum_{k_{m-1}=1}^{k_{m-2}} \frac{q^{k_{m-1}-1}}{(1 - tq^{k_{m-1}-m}) (1 - vq^{k_{m-1}})} \\ & \quad \times \left(\sum_{k_m=1}^{k_{m-1}} \frac{q^{k_m-1}}{1 - vq^{k_m}} \sum_{k_{m-1}=1}^{k_{m-1}-1} \frac{tq^{k_{m-1}-m-1}}{v - tvq^{k_{m-1}-m}} \right). \end{aligned}$$

We perform partial fraction decomposition on

$$F(z) = \frac{(vq; q)_n (tzq^{-m+1}; q)_{m+n-2}}{(vzq; q)_n (tq^{-m} / v; q)_{m-1} (tq^{-m+1}; q)_{m+n-2} (z-1)^{m+1}}.$$

Similarly to the proof of Theorem 2.3, we can prove the above theorem.

Here we omit the proof.

When we set $v = 1$ in Theorem 3.10, we get Theorem 2.3.

Setting $v = q$ in Theorem 3.10, we have

Corollary 3.11. *For $m, n \geq 1$, we have*

$$\begin{aligned} & \sum_{k=1}^n \frac{(q^2; q)_n (q^{m+2} / t; q)_{k-1} (tq^{-2}; q)_{n-k}}{(tq^{-m+1}; q)_{m+n-2} (q; q)_{k-1} (q; q)_{n-k} (1 - q^{k+1})^{m+1}} \left(\frac{t}{q} \right)^{k-1} \\ &= \sum_{k_1=1}^n \frac{q^{k_1-1}}{(1 - tq^{k_1-2}) (1 - q^{k_1+1})} \cdots \sum_{k_{m-1}=1}^{k_{m-2}} \frac{q^{k_{m-1}-1}}{(1 - tq^{k_{m-1}-m}) (1 - q^{k_{m-1}+1})} \\ & \quad \times \left(\sum_{k_m=1}^{k_{m-1}} \frac{q^{k_m-1}}{1 - q^{k_m+1}} - \sum_{k_m=1}^{k_{m-1}-1} \frac{tq^{k_{m-1}-m-2}}{1 - tq^{k_{m-1}-m}} \right). \end{aligned}$$

Letting $n \rightarrow \infty$ in Theorem 3.10, we obtain

Corollary 3.12. *For $m \geq 1$, we have*

$$\begin{aligned} \sum_{k=1}^{\infty} \frac{(vq^{m+1}/t; q)_{k-1}}{(tq^{-m+1}; q)_{m-1}(q; q)_{k-1}(1-vq^k)^{m+1}} \left(\frac{t}{v}\right)^{k-1} &= \frac{(q; q)_{\infty}(t; q)_{\infty}}{(vq; q)_{\infty}(tq^{-1}/v; q)_{\infty}} \\ &= \sum_{k_1=1}^{\infty} \frac{q^{k_1-1}}{(1-tq^{k_1-2})(1-vq^{k_1})} \cdots \sum_{k_{m-1}=1}^{k_{m-2}} \frac{q^{k_{m-1}-1}}{(1-tq^{k_{m-1}-m})(1-vq^{k_{m-1}})} \\ &\quad \times \left(\sum_{k_m=1}^{k_{m-1}} \frac{q^{k_m-1}}{1-vq^{k_m}} \sum_{k_m=1}^{k_{m-1}-1} \frac{tq^{k_m-m-1}}{v-tvq^{k_m-m}} \right). \end{aligned}$$

Setting $m = 1$ in Theorem 3.10, we have the following generalization of the $l = 1$ case of [10, Corollary 3.2].

Corollary 3.13. *For $n \geq 1$, we have*

$$\sum_{k=1}^n \frac{(vq; q)_n (vq^2/t; q)_{k-1} (tq^{-1}/v; q)_{n-k}}{(t, q)_{n-1} (q; q)_{k-1} (q; q)_{n-k} (1-vq^k)^2} \left(\frac{t}{v}\right)^{k-1} = \sum_{k=1}^n \frac{q^{k-1}}{1-vq^k} - \sum_{k=1}^{n-1} \frac{tq^{k-2}}{v-tvq^{k-1}}.$$

4. Two Identities of Fang

We notice that by using partial fraction decomposition technique, we can also give new proofs of two identities found by Fang in [8].

Theorem 4.1 [8, Corollary 3.4]. *We have*

$$\sum_{j=0}^M \begin{bmatrix} M \\ j \end{bmatrix} (-1)^j q^{\binom{j}{2}+2j} \frac{1}{1-zq^j} = \frac{(q; q)_M}{(z; q)_{M+1}} \sum_{j=0}^M (z; q)_j q^j.$$

Proof. Set

$$F(z) = \frac{(q; q)_M}{(z; q)_{M+1}} \sum_{j=0}^M (z; q)_j q^j.$$

Performing partial fraction decomposition on $F(z)$, we have

$$F(z) = \sum_{k=0}^M \frac{b_k}{1-zq^k}. \quad (4.1)$$

For $0 \leq k \leq M$, we have

$$\begin{aligned}
 b_k &= \lim_{z \rightarrow q^{-k}} (1 - zq^k)F(z) \\
 &= \lim_{z \rightarrow q^{-k}} \frac{(q; q)_M}{(z; q)_k (zq^{k+1}; q)_{M-k}} \sum_{j=0}^M (z; q)_j q^j \\
 &= \frac{(q; q)_M}{(q^{-k}; q)_k (q; q)_{M-k}} \sum_{j=0}^M (q^{-k}; q)_j q^j \\
 &= (-1)^k q^{\binom{k+1}{2}} \begin{bmatrix} M \\ k \end{bmatrix} \sum_{j=0}^k (q^{-k}; q)_j q^j \\
 &= \begin{bmatrix} M \\ k \end{bmatrix} (-1)^k q^{\binom{k}{2} + 2k}.
 \end{aligned}$$

The last equation of this chain of equalities follows from the q -Chu-Vandermonde identity [9, Appendix II.6]

$${}_2\phi_1 \left[\begin{matrix} q^{-n}, & a \\ & c \end{matrix} ; q, q \right] = \frac{(c/a; q)_n}{(c; q)_n} a^n, \quad (4.2)$$

by setting $a = q$ and $c = 0$.

According to (4.1), we get Theorem 4.1. □

Theorem 4.2 [8, Corollary 4.1]. *We have*

$$\sum_{j=0}^M \frac{q^j}{(z; q)_{j+1}} = \sum_{j=0}^M \frac{(-1)^j q^{\binom{j}{2} + 2j}}{(1 - zq^j)(q; q)_j (q; q)_{M-j}}.$$

Proof. Set

$$F(z) = \sum_{j=0}^M \frac{q^j}{(z; q)_{j+1}}.$$

Performing partial fraction decomposition on $F(z)$, we have

$$F(z) = \sum_{k=0}^M \frac{b_k}{1 - zq^k}. \quad (4.3)$$

For $0 \leq k \leq M$, we have

$$\begin{aligned} b_k &= \lim_{z \rightarrow q^{-k}} (1 - zq^k) F(z) \\ &= \lim_{z \rightarrow q^{-k}} (1 - zq^k) \sum_{j=0}^M \frac{q^j}{(z; q)_{j+1}} \\ &= \lim_{z \rightarrow q^{-k}} (1 - zq^k) \sum_{j=k}^M \frac{q^j}{(z; q)_{j+1}} \\ &= \lim_{z \rightarrow q^{-k}} \sum_{j=k}^M \frac{q^j}{(z; q)_k (zq^{k+1}; q)_{j-k}} \\ &= \sum_{j=k}^M \frac{q^j}{(q^{-k}; q)_k (q; q)_{j-k}} \\ &= \frac{(-1)^k q^{\binom{k}{2} + 2k}}{(q; q)_k} \sum_{j=0}^{M-k} \frac{q^j}{(q; q)_j} \\ &= \frac{(-1)^k q^{\binom{k}{2} + 2k}}{(q; q)_k (q; q)_{M-k}}. \end{aligned}$$

The last equation follows from the q -Chu-Vandermonde identity (4.2) by setting $a = 0$ and $c = q^{-n}$.

According to (4.3), we complete the proof of Theorem 4.2. $\square$

References

- [1] W. Chu and A. Marini, Partial fractions and trigonometric identities, *Adv. in Appl. Math.* 23 (1999), 115-175.
- [2] W. Chu, A binomial coefficient identity associated with Beukers' conjecture on Apéry numbers, *Electron J. Combin.* 11 (2004), Note 15, 3 pp.
- [3] W. Chu, Partial fraction decompositions and trigonometric sum identities, *Proc. Amer. Math. Soc.* 136 (2008), 229-237.
- [4] W. Chu and Y. You, Binomial symmetries inspired by Bruckman's problem, *Filomat* 24 (2010), 41-46.
- [5] W. Chu, C. Y. Wang and W. L. Zhang, Partial fractions and q -binomial determinant identities *CUBO* 12 (2010), 1-12.
- [6] S. Corteel and J. Lovejoy, Overpartitions, *Trans. Amer. Math. Soc.* 356 (2004), 1623-1635.
- [7] K. Dilcher, Some q -series identities related to divisor functions, *Discrete Math.* 145 (1995), 83-93.
- [8] J. P. Fang, Some applications of q -differential operator, *J. Korean Math. Soc.* 47 (2010), 223-233.
- [9] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, 2nd Edition *Encyclopedia of Mathematics and its Applications*, Vol. 96, Cambridge University Press, Cambridge, 2004.
- [10] V. J. W. Guo and C. Zhang, Some further q -series identities related to divisor functions, *Ramanujan J.* 25 (2011), 295-306.
- [11] H. Prodinger, Some applications of the q -Rice formula, *Random Struct. Algorithms* 19 (2001), 552-557.

